

7. Wolfe JH. Endovascular repair (EVAR) of abdominal aortic aneurysms: the case against. *Ann R Coll Surg Engl* 2004;86:379-80.
8. EVAR trial participants. Endovascular aneurysm repair versus open repair in patients with abdominal aortic aneurysm (EVAR trial 1): randomised controlled trial. *Lancet* 2005;365(9478):2179-86.
9. Blankensteijn JD, de Jong SE, Prinssen M, van der Ham AC, Buth J, van Sterkenburg SM, Verhagen HJ, Buskens E, Grobbee DE. Two-year outcomes after conventional or endovascular repair of abdominal aortic aneurysms. *N Engl J Med* 2005 Jun 9;352(23):2443-5.
10. Elkouri S, Martelli E, Głowiczki P, McKusick MA, Panneton JM, Andrews JC, Noel AA, Bower TC, Sullivan TM, Rowland C, Hoskin TL, Cherry KJ. Most patients with abdominal aortic aneurysm are not suitable for endovascular repair using currently approved bifurcated stent-grafts. *Vasc Endovasc Surg* 2004;38(5):401-12.
11. Hinchliffe RJ, Goldberg J, Macsweeney STR and on behalf of the Zenith Group. A UK multi-centre experience with a second generation endovascular stent graft: results from the Zenith users group. *Eur J Vasc Surg* 2004;27(1):51-5.
12. Ouriel K, Clair DG, Greenberg RK, Lyden SP, O'Hara PJ, Sarac TP, Srivastava SD, Butler B, Sampram ES. Endovascular repair of abdominal aortic aneurysms: device specific outcome. *J Vasc Surg* 2003;37:991-8.
13. Laheij RJF, Buth J, Harris PL, Moll FL, Stelter WJ, Verhoeven ELG. Need for secondary interventions after endovascular repair of abdominal aortic aneurysms. Intermediate-term follow-up results of a European collaborative registry (EUROSTAR). *Br J Surg* 2000;87:1666-73.
14. Peppelenbosh N, Buth J, Harris PL, Marrewijk C, Fransen G, for the Eurostar collaborators. Diameter of abdominal aortic aneurysm and outcome of endovascular aneurysm repair: does size matter? A report from EUROSTAR. *J Vasc Surg* 2004;39:288-97.

Submitted Jul 3, 2005; accepted Sep 26, 2005.

INVITED COMMENTARY

J. Adam van der Vliet, MD, PhD, and Jan D. Blankensteijn, MD, PhD, Nijmegen, The Netherlands

The authors compared the suitability for endovascular repair (EVAR) in patients with incidentally diagnosed and screen-detected aneurysms. In view of the potential effects on the future health economics of aneurysm screening and EVAR, this is a topic of considerable interest. The study was initiated after the authors' observation that screen-detected aneurysms tended to have longer infrarenal necks. However, early abdominal aortic aneurysm (AAA) detection did not lead to an increase in suitability for EVAR in this study.

Before the outcome of this study is accepted, several issues must be addressed. First, the study lacks a proper power analysis, and with recruitment arms of 41 and 31 patients, a type II error is imminent. One may further question the efficacy of the ultrasound screening program to which the study population was exposed, because the median aneurysm diameter upon referral was 65 mm (range, 51-79 mm), after a mean of nine surveillance scans per patient, typically spanning a minimum of 2 years. As a result, the screen-detected aneurysms were relatively large at the time of referral, whereas EVAR suitability is expected to be higher in smaller AAAs. The study was conducted in a vascular center where EVAR is not offered as a treatment option, and all patients underwent open repair. Presumably, a center with little or no EVAR

experience will tend to use rather rigid criteria for this treatment option as compared with EVAR centers. Common indications for surgery were applied in this study: aneurysm diameter larger than 55 mm, rapid expansion of more than 10 mm/y, or a tender aneurysm. This is in accordance with most EVAR studies. The determination of suitability for EVAR, however, depended on the—rather poorly defined—manufacturer's criteria for a single device, whereas in most endovascular programs, EVAR eligibility is determined on the basis of the availability of multiple devices. In addition, suitability for EVAR is never an objective qualification, irrespective of manufacturer's guidelines, which are commonly rather conservative (for understandable reasons). Many clinicians, for instance, consider neck thrombus not to be a contraindication to EVAR, although this was the second most common exclusion criterion in this study. This may be an explanation for the low rates of suitability for EVAR of 41% and 45% in the screen-detected and incidentally diagnosed AAAs, respectively, whereas in many centers 60% of AAA patients receive an endograft. It is quite possible that with more liberal criteria for EVAR, screen-detected aneurysms would have shown a higher suitability rate than the incidentally diagnosed aneurysms. Caution is therefore appropriate in the interpretation and generalization of the outcome of this study.